



HELIOS · A

HELIOS-A: Study of Vutrisiran in Patients with hATTR Amyloidosis

David Adams¹, Ivailo L Tournev^{2,3}, Mark S Taylor⁴, Teresa Coelho⁵, Violaine Planté-Bordeneuve⁶, John L Berk⁷, Alejandra González-Duarte⁸, Julian D Gillmore⁹, Soon-Chai Low¹⁰, Yoshiki Sekijima¹¹, Laura Obici¹², Chongshu Chen¹³, Prajakta Badri¹³, Seth Arum¹³, John Vest¹³, Michael Polydefkis¹⁴

¹Neurology Department, APHP, CHU Bicêtre, INSERM U1195, Université Paris-Saclay, ²Clinic of Nervous Diseases, University Hospital Aleksandrovska, Department of Neurology, Medical University – Sofia, ³Department of Cognitive Sciences, New Bulgarian University, ⁴Department of Clinical Immunology and Allergy, Westmead Hospital and Westmead Clinical School, University of Sydney, ⁵Hospital de Santo António, Centro Hospitalar do Porto, ⁶Neurology – Amyloid network, CHU Henri Mondor – Assistance Publique Hopitaux de Paris, ⁷Boston Medical Center, ⁸Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, ⁹National Amyloidosis Centre, University College London, Royal Free Hospital, ¹⁰Division of Neurology, Department of Medicine, University of Malaya, ¹¹Department of Medicine (Neurology & Rheumatology), Shinshu University School of Medicine, ¹²Amyloidosis Research and Treatment Centre, IRCCS Fondazione Policlinico San Matteo, ¹³Alnylam Pharmaceuticals, ¹⁴Department of Neurology, Johns Hopkins University School of Medicine

January 21–22, 2022 || 26th édition of the Journées de la Société Francophone du Nerf Périphérique (SFNP)

Background and Rationale

hATTR Amyloidosis, Also Known as ATTRv Amyloidosis

- Rare, underdiagnosed, inherited, rapidly progressive, debilitating, and fatal disease
- Caused by variants in the *TTR* gene that result in misfolded TTR accumulating as amyloid deposits in multiple organs and tissues¹⁻⁴
- The majority of individuals develop a mixed phenotype of polyneuropathy and cardiomyopathy^{5,6}


Vutrisiran

- Investigational, subcutaneously administered RNAi therapeutic targeting hepatic production of variant and wt TTR in development for the treatment of ATTR amyloidosis^{7,8}

Patisiran

- RNAi therapeutic administered via IV infusion, approved for the treatment of the polyneuropathy of hATTR amyloidosis based on the Phase 3, placebo-controlled APOLLO trial⁹⁻¹²

Therapeutic Hypothesis



ATTRv, hereditary transthyretin (v for variant); ESC, enhanced stabilization chemistry; GalNAc, *N*-acetylgalactosamine; hATTR, hereditary transthyretin-mediated; IV, intravenous; Q3M, every 3 months; RNAi, ribonucleic acid interference; SC, subcutaneous; TTR, transthyretin; wt, wild-type.


1. Hanna M. *Curr Heart Fail Rep* 2014;11:50-57; 2. Hawkins PN et al. *Ann Med* 2015;47:625-638; 3. Damy T et al. *J Cardiovasc Transl Res* 2015;8:117-127; 4. Mohty D et al. *Arch Cardiovasc Dis* 2013;106:528-540; 5. Rapezzi C et al. *Eur Heart J* 2013;34:520-528; 6. Coelho T et al. *Curr Med Res Opin* 2013;29:63-76; 7. Habtemariam BA et al. *Clin Pharmacol Ther* 2021;109:372-382; 8. Nair JK et al. *J Am Chem Soc* 2014;136:16958-16961; 9. Alnylam Pharmaceuticals. US prescribing information: ONPATTRO® (patisiran) lipid complex injection, for intravenous use. February 2020; 10. Adams D et al. *N Engl J Med* 2018;379:11-21; 11. APOLLO: NCT01960348; 12. Alnylam France, Résumé des caractéristiques du produit ONPATTRO® (patisiran).

Vutrisiran Phase 3 HELIOS·A Study in Patients with Hereditary Transthyretin-Mediated Amyloidosis Polyneuropathy



Month 18 Results

- As previously reported, the primary endpoint of change from baseline in mNIS+7 at Month 9 was met¹



^aHigher scores of mNIS+7 indicate more neurologic impairment (range, 0 to 304). ^bHigher scores of Norfolk QOL-DN indicate worse quality of life (range, -4 to 136). ^c10-MWT speed (m/s) = 10 meters/mean time (seconds) taken to complete two assessments at each visit, imputed as 0 for patients unable to perform the walk; lower speeds indicate worse ambulatory function. ^dLower scores of mBMI (weight [in kg/m²] × serum albumin [in g/L]) indicate worse nutritional status. ^eLower scores of R-ODS indicate more disability (range, 0 to 48). ^fTc scintigraphy was only performed at select sites.

10-MWT, 10-meter walk test; hATTR, hereditary transthyretin-mediated amyloidosis; IV, intravenous; KPS, Karnofsky performance status; mBMI, modified body mass index; mNIS+7, modified Neuropathy Impairment Score +7; NIS, Neuropathy Impairment Score; Norfolk QOL-DN, Norfolk Quality of Life-Diabetic Neuropathy; PND, polyneuropathy disability; Q3M, every 3 months; Q3W, every 3 weeks; R-ODS, Rasch-built Overall Disability Scale; SC, subcutaneous; Tc, technetium; TTR, transthyretin.

1. Adams D et al. *Neurology* 2021;96(15 Supplement):1234.



Baseline Demographic and Disease Characteristics

Characteristic	APOLLO	HELIOS-A	
	Placebo (n=77)	Vutrisiran (n=122)	Patisiran (n=42)
Age (years), median (range)	63 (34, 80)	60 (26, 85)	60 (31, 81)
Males, n (%)	58 (75)	79 (65)	27 (64)
TTR genotype, n (%)			
V30M	40 (52)	54 (44)	20 (48)
Non-V30M	37 (48)	68 (56)	22 (52)
NIS, mean (range)	57 (7, 126)	43 (5, 127)	43 (6, 116)
Previous tetramer stabilizer use, n (%)	41 (53.2)	75 (61.5)	33 (78.6)
PND score ^a , n (%)			
I: preserved walking, sensory disturbances	20 (26)	44 (36)	15 (36)
II: impaired walking but can walk without stick or crutch	23 (30)	50 (41)	17 (40)
IIIA: walk with 1 stick or crutch	22 (29)	16 (13)	7 (17)
IIIB: walk with 2 sticks or crutches	11 (14)	12 (10)	3 (7)
Cardiac subpopulation, n (%) ^{b,c}	36 (47)	40 (33)	14 (33)

^aOne patient (1.3%) in the external placebo group had a PND score of IV defined as confined to wheelchair or bedridden (not shown on the slide). ^bCardiac subpopulation was defined as patients who had pre-existing evidence of cardiac amyloid involvement (baseline LV wall thickness ≥ 1.3 cm and no aortic valve disease or hypertension in medical history). ^cSelect echocardiogram parameters were reread for the Month 18 analysis and the cardiac subpopulation was rederived based on baseline LV wall thickness values after the re-read. As a result, in the Month 18 analysis the cardiac subpopulation status of 9 patients receiving vutrisiran was reclassified and 1 patient receiving patisiran was added to the cardiac subpopulation compared with the cardiac subpopulation defined in the Month 9 analysis.

Statistically Significant Improvement in Neuropathy Impairment and Quality of Life with Vutrisiran vs External Placebo at Month 18


- Improvement was observed across all prespecified patient subgroups, components, and subdomains of mNIS+7 and Norfolk QOL-DN (data not shown)
- Improvement relative to baseline^a in mNIS+7 (48.3% [vutrisiran] vs 3.9% [placebo]) and Norfolk QOL-DN (56.8% vs 10.4%)**
- Consistent treatment effects in vutrisiran and patisiran groups in HELIOS-A (data not shown)



^aImprovement defined as patients with <0-point increase from baseline to 18 months. ^bmITT population (all randomized patients who received any amount of study drug). Value of n is the number of evaluable patients at each timepoint. Data plotted for mNIS+7 and Norfolk QOL-DN at Month 9 are ANCOVA/multiple imputation model data and data plotted at Month 18 are MMRM model data. ^cAt baseline, the mean (±SD) mNIS+7 was 60.6 (36.0) in the vutrisiran group and 74.6 (37.0) in the external placebo group. ^dAt baseline, the mean (±SD) Norfolk QOL-DN score was 47.1 (26.3) in the vutrisiran group and 55.5 (24.3) in the external placebo group.

ANCOVA, analysis of covariance; CI, confidence interval; LS, least squares; LSMD, LS mean difference; mITT, modified intent-to-treat; MMRM, mixed model for repeated measures; mNIS+7, modified Neuropathy Impairment Score +7; Norfolk QOL-DN, Norfolk Quality of Life-Diabetic Neuropathy; SD, standard deviation; SEM, standard error of the mean.

Statistically Significant Improvement in Secondary Endpoints with Vutrisiran vs External Placebo at Month 18




^amITT population (all randomized patients who received any amount of study drug) for all endpoints. Value of n is the number of evaluable patients at each timepoint. Data plotted for 10-MWT, mBMI and R-ODS at Month 9 are ANCOVA/multiple imputation model data and data plotted at Month 18 are MMRM model data. ^bAt baseline, the mean (±SD) 10-MWT was 1.006 (0.393) in the vutrisiran group and 0.790 (0.319) in the external placebo group. ^cAt baseline, the mean (±SD) mBMI was 1057.4 (233.8) in the vutrisiran group and 989.9 (214.2) in the external placebo group. ^dAt baseline, the mean (±SD) R-ODS was 34.1 (11.0) in the vutrisiran group and 29.8 (10.8) in the external placebo group.

10-MWT, 10-meter walk test; ANCOVA, analysis of covariance; CI, confidence interval; LS, least squares; LSMD, LS mean difference; mBMI, modified body mass index; mITT, modified intent-to-treat; MMRM, mixed model repeated measures; R-ODS, Rasch-built Overall Disability Scale; SD, standard deviation; SE, standard error.

Improvement in Exploratory Assessment of NT-proBNP with Vutrisiran vs External Placebo at Month 18

Change from Baseline in NT-proBNP (mITT Population)^a






^aNT-proBNP is a measure of cardiac stress, with higher values indicating a greater level of cardiac stress. ^bAt baseline, NT-proBNP geometric mean (SE) was 273.0 (42.2) ng/L in the vutrisiran group (n=122) and 531.3 (86.7) ng/L in the APOLLO placebo group (n=75). Number of evaluable patients at each timepoint are shown. Data plotted for NT-proBNP at Month 9 are ANCOVA/multiple imputation model data and data plotted at Month 18 are MMRM model data. ANCOVA, analysis of covariance; mITT, modified intent-to-treat; MMRM, mixed model repeated measures; NT-proBNP, N-terminal pro-brain natriuretic peptide; SE, standard error.

Assessments of Cardiac Amyloid Involvement with Vutrisiran

- Echocardiographic parameters at Month 18 trended toward improvement with vutrisiran compared with the external placebo group (exploratory endpoints)

- Cardiac uptake of ^{99m}Tc on scintigraphy imaging at Month 18 was reduced with vutrisiran compared with baseline in a planned cohort (exploratory endpoints)




^amITT population. ^bPatients from a planned cohort of the mITT population for whom the relevant 18-month data were available. ^cImproved: <0 increase from baseline. ^dNot improved: ≥0 increase from baseline.

LS, least squares; LV, left ventricular; mITT, modified intent-to-treat; SE, standard error; Tc, technetium.

Rapid and Sustained Reduction in Serum TTR Levels with Vutrisiran

- Vutrisiran achieved a mean steady-state serum TTR reduction from baseline of 88% (SD: 16%)
- TTR reduction with vutrisiran was non-inferior to that observed with the within-study patisiran reference comparator (secondary endpoint) over 18 months^a

Percent Change from Baseline in Serum TTR Levels




N evaluable																		
Vutrisiran (n = 122)	122	114	109	119	106	117	92	118	115	56	116	42	118	15	118	100	114	98
Patisiran (n = 42)	42	42	41	41	37	38	39	34	39	23	40	23	36	9	37	36	38	32

^aAs assessed by mean trough serum TTR levels.
SD, standard deviation; SE, standard error; TTR, transthyretin.


Month 18 HELIOS-A Vutrisiran Efficacy Results Consistent with APOLLO Patisiran

Findings Consistent with Similar Serum TTR Reduction Seen with Vutrisiran and Patisiran

Vutrisiran Efficacy^a vs Placebo
Standardized Effect Sizes from HELIOS-A



Patisiran Efficacy^b vs Placebo
Standardized Effect Sizes from APOLLO



^aHELIOS-A mITT population. ^bAPOLLO mITT population.

10-MWT, 10-meter walk test; LV, left ventricular; mBMI, modified body mass index; mITT, modified intent-to-treat; mNIS+7, modified Neuropathy Impairment Score +7; Norfolk QOL-DN, Norfolk Quality of Life-Diabetic Neuropathy; NT-proBNP, N-terminal pro-brain natriuretic peptide; R-ODS, Rasch-built Overall Disability Scale.

HELIOS-A Safety Summary^a

The majority of AEs were mild or moderate in severity

- No drug-related discontinuations or deaths
- Three study discontinuations (2.5%) due to AEs in the vutrisiran arm (two due to death, as previously reported; one due to heart failure), none of which were considered related to study drug
 - One death due to COVID-19 pneumonia and the other due to iliac artery occlusion
- As previously reported, two SAEs deemed related to vutrisiran by investigators:
 - Dyslipidemia and urinary tract infection
- AEs ≥10% in the vutrisiran group included fall, pain in extremity, diarrhea, peripheral edema, urinary tract infection, arthralgia, and dizziness
- Injection-site reactions were reported in 5 patients (4.1%) receiving vutrisiran; all were mild and transient
- No safety signals regarding liver function tests, hematology, or renal function related to vutrisiran

HELIOS-A Safety Summary^a

At least one event, n (%)	APOLLO	HELIOS-A	
	Placebo (n=77)	Vutrisiran (n=122)	Patisiran (n=42)
AEs	75 (97.4)	119 (97.5)	41 (97.6)
SAEs	31 (40.3)	32 (26.2)	18 (42.9)
Severe AEs	28 (36.4)	19 (15.6)	16 (38.1)
AEs leading to treatment discontinuation	11 (14.3)	3 (2.5)	3 (7.1)
AEs leading to stopping study participation	9 (11.7)	3 (2.5)	2 (4.8)
Deaths	6 (7.8)	2 (1.6)	3 (7.1)

^aData reported during 18-month treatment period.
AE, adverse event; SAE, serious AE.

Summary

- As previously reported, vutrisiran met the HELIOS-A primary endpoint (mNIS+7) at 9 months¹
- Vutrisiran met all 18-month secondary endpoints
 - Maintained statistically significant improvement in mNIS+7 compared with external placebo
 - Improvement in QOL (Norfolk QOL-DN), gait speed (10-MWT), nutritional status (mBMI), and disability (R-ODS), compared with external placebo
 - Robust and sustained TTR reduction, non-inferior to within-study patisiran
- Improvements in certain exploratory cardiac measures, including NT-proBNP, compared with external placebo
- Vutrisiran had an acceptable safety profile
 - The majority of AEs were mild or moderate in severity with the most common treatment-related AEs being injection site reactions
- HELIOS-A continues to investigate the efficacy and safety of vutrisiran through an ongoing extension period

10-MWT, 10-meter walk test; - AE, adverse event; mBMI, modified body mass index; mNIS+7, modified Neuropathy Impairment Score +7; Norfolk QOL-DN, Norfolk Quality of Life-Diabetic Neuropathy; NT-proBNP, N-terminal pro-brain natriuretic peptide; QOL, quality of life; R-ODS, Rasch-built Overall Disability Scale; TTR, transthyretin.

1. Adams D et al. *Neurology* 2021;96(15 Supplement):1234.



Thank you to the patients, their families,
investigators, study staff, and collaborators for
their participation in the **HELIOS-A study**