

Vutrisiran: Post-Orthotopic Liver Transplant

The following information is provided in response to your unsolicited inquiry. It is intended to provide you with a review of the available scientific literature and to assist you in forming your own conclusions in order to make healthcare decisions. This document is not for further dissemination or publication without authorization.

The full Prescribing Information for AMVUTTRA® (vutrisiran) is provided [here](#). Alnylam Pharmaceuticals does not recommend the use of its products in any manner that is inconsistent with the approved Prescribing Information. This resource may contain information that is not in the approved Prescribing Information.

If you are seeking additional scientific information related to Alnylam medicines, you may visit the Alnylam US Medical Affairs website at RNAiScience.com.

SUMMARY

- In the HELIOS-A and HELIOS-B studies, patients were excluded if they had a liver transplant or were likely to undergo liver transplantation during the study period.^{1,2}

INDEX

[Clinical Data](#) – [Label Information](#) – [Abbreviations](#) – [References](#)

CLINICAL DATA

Phase 3 HELIOS-A Study

HELIOS-A was a phase 3, global, randomized, open-label study designed to evaluate the efficacy and safety of vutrisiran in patients with hATTR-PN. Patients were randomized (3:1) to receive either vutrisiran 25 mg every 3 months by subcutaneous injection (n=122) or patisiran 0.3 mg/kg every 3 weeks by IV infusion (as a reference group, n=42) for 18 months. This study used the placebo arm of the APOLLO study as an external control arm (n=77) for the primary endpoint and most other efficacy endpoints. The primary endpoint was the change from baseline in mNIS+7 at 9 months.¹

Exclusion Criteria

Patients were excluded from the study if any of the following criteria applied¹:

- Had a liver transplant or were likely to undergo liver transplantation during the 18-month treatment period of the study

Phase 3 HELIOS-B Study

HELIOS-B was a phase 3, global, randomized, double-blind, placebo-controlled, multicenter study designed to evaluate the efficacy and safety of vutrisiran in patients with ATTR-CM, including both hATTR and wtATTR. Patients were randomized (1:1) to receive either vutrisiran 25 mg (n=326) or placebo (n=329) every 3 months by subcutaneous injection for up to 36 months. The primary endpoint was the composite endpoint of all-cause mortality and recurrent CV events (CV hospitalizations and urgent heart failure visits) at the end of the double-blind exposure period in the overall population and in the vutrisiran monotherapy population (patients not receiving tafamidis at baseline).³

Exclusion Criteria

Patients were excluded from the study if any of the following criteria applied²:

- Prior or anticipated (during the first 12 months after randomization) heart, liver, or other organ transplant or implantation of left-ventricular assist device

AMVUTTRA PRESCRIBING INFORMATION – RELEVANT CONTENT

The CLINICAL PHARMACOLOGY section provides the following information⁴:

Specific Populations

No clinically significant differences in the pharmacokinetics of vutrisiran were observed based on age, sex, race, mild and moderate renal impairment (eGFR ≥ 30 to < 90 mL/min/1.73 m²), or mild hepatic impairment (total bilirubin $\leq 1 \times$ ULN and AST $> 1 \times$ ULN, or total bilirubin > 1.0 to $1.5 \times$ ULN and any AST). Vutrisiran has not been studied in patients with severe renal impairment, end-stage renal disease, moderate or severe hepatic impairment, or in patients with prior liver transplant.

ABBREVIATIONS

AST = aspartate aminotransferase; ATTR-CM = transthyretin amyloidosis with cardiomyopathy; CV = cardiovascular; eGFR = estimated glomerular filtration rate; hATTR = hereditary transthyretin amyloidosis; hATTR-PN = hereditary transthyretin amyloidosis with polyneuropathy; IV = intravenous; mNIS+7 = modified neuropathy impairment score +7; ULN = upper limit of normal; wtATTR = wild-type transthyretin amyloidosis.

Updated 17 Sept 2024

REFERENCES

1. Adams D, Tournev IL, Taylor MS, et al. Efficacy and safety of vutrisiran for patients with hereditary transthyretin-mediated amyloidosis with polyneuropathy: a randomized clinical trial. *Amyloid*. 2023;30(1):18-26. doi:10.1080/13506129.2022.2091985
2. Protocol for: Fontana M, Berk JL, Gillmore JD, et al. Vutrisiran in patients with transthyretin amyloidosis with cardiomyopathy. *N Engl J Med*. 2024. doi:10.1056/NEJMoa2409134
3. Fontana M, Berk JL, Gillmore JD, et al. Vutrisiran in patients with transthyretin amyloidosis with cardiomyopathy. *N Engl J Med*. August 2024. doi:10.1056/NEJMoa2409134
4. AMVUTTRA (vutrisiran) Prescribing Information. Cambridge, MA: Alnylam Pharmaceuticals, Inc.